

Effect of power density on  
sensitised human RBC in  
PBS/Mg/Glucose




Fig.1

Fig 2A





Fig 2B






Fig. 3

Ultrasound sensitivity during  
storage at 4°C in plasma




Fig. 4

5/20




Fig. 5




Fig. 6

7/20

Fig 7A




Fig 7B





Fig 7C



8/20




Fig. 8

002221 122200 09748063






Fig. 9




Fig.10





Fig.11

12/20

+U/S



-U/S




Fig.12

00222T E 908460

Fig. 13A RbRBC + Electrosensitisation




Fig.13B RbRBC unmodified





Fig.13C RbRBC plus 2.5% glutaraldehyde



002227 E9084650




Fig.13D

09748063.122200




Fig. 14

16/20

# *in vivo* survival of modified erythrocytes in rabbit




Fig.15



Survival of loaded and  
sensitised rabbit erythrocytes  
in circulation




Fig.16

**CIRCULATING PHANTOM**  
**Ultrasound-induced antibody**  
**payload from spiked whole blood**  
**at 40% haematocrit**  
**(US - continuous wave, 5W/cm<sup>2</sup>)**




Fig.17

Clearance of anti-Hu IgG from rabbit  
circulation (n=3) as measured by  
ELISA




Fig. 18

## Ultrasound-mediated release of antibody payload *in vivo*




Fig. 19